

Phenotypic Variation of *Helicobacter pylori* Isolates from Geographically Distinct Regions Detected by Lectin Typing

Sean O. Hynes,^{1,2} Nathalie Broutet,³ Eurohepygast Study Group,^{3†} Torkel Wadström,² Marika Mikelsaar,⁴ Paul W. O'Toole,⁵ John Telford,⁶ Lars Engstrand,⁷ Shigeru Kamiya,⁸ Andreas F. Mentis,⁹ and Anthony P. Moran^{1*}

Department of Microbiology, National University of Ireland, Galway, Ireland¹; Department of Medical Microbiology, Dermatology and Infection, University of Lund, Lund,² and Swedish Institute for Infectious Disease Control, Solna,⁷ Sweden; Epidemiology Unit of Digestive Diseases, Laboratory of Bacteriology, University of Victor Segalen Bordeaux 2, Bordeaux, France³; Department of Medical Microbiology, University of Tartu, Tartu, Estonia⁴; Institute of Molecular BioSciences, Massey University, Palmerston North, New Zealand⁵; Department of Molecular Biology, Chiron Vaccines, Immunobiological Research Institute, Siena, Italy⁶; Department of Infectious Diseases, Kyorin University School of Medicine, Tokyo, Japan⁸; and Laboratory of Bacteriology, Hellenic Pasteur Institute, Athens, Greece⁹

Received 31 May 2001/Returned for modification 15 August 2001/Accepted 15 October 2001

A total of 309 *Helicobacter pylori* isolates from 18 different countries were analyzed with a previously developed lectin typing system. The system was developed by using a proteolytic pretreatment to enhance the carbohydrate fraction of the sample. Four lectins from *Ulex europaeus*, *Lotus tetragonolobus*, *Erythrina cristigali*, and *Triticum vulgare* were used to type the strains. The lectins were chosen for their specificities for sugars commonly encountered in the lipopolysaccharide of *H. pylori*. The isolates were received from their parent institutions as pellets of biomass and were typed at one of three centers (in Ireland, Sweden, and Estonia). All 16 possible lectin reaction patterns were observed in the study, with the isolates with the predominant pattern exhibiting reactions with all the lectins in the panel. For European patients suffering from gastritis, an association was noted between lectin reaction pattern MH4 and atrophic chronic gastritis; isolates with lectin reaction pattern MH4 were isolated from patients with atrophic chronic gastritis, whereas isolates with this pattern were not isolated from patients with chronic gastritis ($P = 0.0006$). In addition, statistically significant relationships were noted between the lectin reaction pattern and the associated pathology of isolates from the Swedish population. Isolates with patterns MH13 and MH16, which had low lectin reactivities, correlated with nonulcer disease ($P = 0.0025$ and $P = 0.0002$, respectively), and all four isolates from adenocarcinoma patients were characterized as possessing reaction pattern MH16. In contrast, isolates with lectin reaction patterns MH1 and MH10, which had high lectin reactivities, were associated with ulcer disease ($P = 0.046$ and $P = 0.0022$, respectively).

Infection with *Helicobacter pylori* is associated with a number of pathologies, including chronic gastritis, peptic ulcer disease, gastric adenocarcinoma, and gastric mucosa-associated lymphoid tissue lymphoma (9). The reason for these differences in clinical outcomes associated with *H. pylori* infection and the reason why some infected individuals remain asymptomatic are not known. It has been suggested that host or environmental factors may play a role in these differences or that certain more virulent strains of *H. pylori* exist (6, 29).

Numerous typing systems have been used to differentiate and group strains of *H. pylori*. The majority of these procedures have involved genotypic typing methods, which have noted differences between strains of *H. pylori* isolated from distinct host populations (4, 14, 18, 25, 31). The results of these studies strongly suggest that host-adapted strains exist. In addition, genotypic characteristics such as the presence of the *cag* pathogenicity island and the presence of *vacA* type s1a and expres-

sion of the cytotoxin-associated gene A product (CagA) or vacuolating toxin (VacA) have been shown to be markers for specific pathologies in certain populations (3, 24, 29, 32). However, such associations have not been found for all populations (12, 17).

A number of phenotypic typing systems for *H. pylori*, including serotyping, are based upon variation in the O side chain of lipopolysaccharide (LPS) of the bacterium, especially the expression of Lewis (Le) blood group determinants (19, 26). Several serological studies have suggested that Le antigens may act as markers for strains associated with specific pathologies, in particular, ulcer disease (8, 17, 32; N. Broutet, XIIIth Int. Workshop Gastrointestinal Pathology and *Helicobacter pylori*, Gut 47[Suppl. I]:A27–A28, 2000). Previously, a novel phenotypic typing system based on lectin typing which can be successfully applied to *H. pylori* was developed (10). This system has advantages over serotyping, particularly that with anti-Le monoclonal antibodies. First, since antibodies for a conserved set of structures are used, such serotyping may exhibit a low level of discriminatory ability; second, a large number (15%) of untypeable strains occur (26); third, technical expertise is required for the performance of the tests (19, 26); and fourth, inconsistencies between test systems that use anti-Le antibodies in different formats have been demonstrated (1, 11).

* Corresponding author. Mailing address: Department of Microbiology, National University of Ireland, Galway, Ireland. Phone: 353-91-524411, ext. 3163. Fax: 353-91-525700. E-mail: anthony.moran@nuigalway.ie.

† See the Appendix for the members of the Eurohepygast Study Group.

Moreover, the lectin typing system does not require the raising of antibodies in animals, as is required for serotyping, and is ethically sound, less time-consuming, and economically advantageous (10). Lectins have been proposed for use as diagnostic and typing tools in a number of studies for other bacteria (5, 13, 23, 34). The present system was developed to detect variations in carbohydrate expression by *H. pylori* LPS. Moreover, the use of lectin-blotting analyses in a previous study (10) established that lectins in the typing system bind predominantly with carbohydrates in the accessible O-side-chain region of LPS but reactions with carbohydrates in the LPS core were not excluded. The system has proved reproducible, sensitive, and easy to use when applied to a group of strains isolated from a geographically distinct population in Ireland (10). The present study reports on the application of this typing system to a large number of isolates from different geographical locations worldwide. The aim of our study was to examine whether the host population or associated gastroduodenal pathology was reflected in phenotypic differences between strains of *H. pylori* as detected by the lectin typing system.

MATERIALS AND METHODS

Origins of bacterial samples. *H. pylori* was isolated from antral biopsy material from 309 patients presenting at open-access endoscopy units and undergoing upper gastrointestinal endoscopy. The patients were from 18 different countries, and each patient presented in his or her home country with dyspeptic symptoms. The patients exhibited a variety of endoscopy-proven disorders, including gastritis, peptic ulcer disease, atrophic chronic gastritis (ACG) with or without intestinal metaplasia (IM) (ACG \pm IM), and gastric adenocarcinoma. Patients presenting with gastritis were classified by using the Sydney system (27). For inclusion in the European cohort (Eurohepygast Study Group) of ACG \pm IM patients, patients had to present themselves in one of the participating centers, to be between 18 and 75 years of age, and to have had an endoscopy in which a biopsy diagnosis of chronic gastritis was made. In addition, a small proportion (3%) of the bacterial isolates were obtained from asymptomatic patients. The country or population of origin and the number of the total of 309 *H. pylori* isolates from each country were as follows: Belgium, $n = 1$; Bulgaria, $n = 17$; China, $n = 12$; Czech Republic, $n = 8$; Estonia, $n = 28$; Finland, $n = 4$; France, $n = 18$; Germany, $n = 9$; Greece, $n = 35$; Hungary, $n = 1$; Ireland, $n = 13$; Italy, $n = 4$; Japan, $n = 20$; the Polynesian population of New Zealand, $n = 10$; Poland, $n = 13$; Portugal, $n = 16$; Spain, $n = 19$; and Sweden, $n = 81$.

Bacteria and culture conditions. The bacteria were routinely grown on 5% blood agar under microaerobic conditions with 10% CO₂ and 5% O₂, and the purity of the cultures was established by the parent laboratory by using previously defined criteria (16). The bacteria were subsequently transported as frozen pellets of biomass which had been washed once by suspension in 0.1 M phosphate-buffered saline (PBS; pH 7.2) with subsequent centrifugation at 3,000 \times g for 10 min. In addition to pellets of biomass, some isolates were received as frozen stock cultures in Trypticase soy broth with 10% (vol/vol) glycerol and were subsequently cultured as described above, and biomass was obtained for comparative typing with transported pellets.

Treatment of bacterial samples prior to lectin typing. Bacterial cells were subjected to proteolytic degradation before use in the lectin typing assay described previously (10). Briefly, stored samples were washed once in PBS and were then resuspended in 5 ml of PBS adjusted to pH 4 for 30 min to induce autolysis of the cells and protein release. Subsequently, the cells were washed twice in PBS. The resultant pellet was resuspended in 5 ml of PBS containing 0.1 mg of proteinase K (Sigma Chemical Co., St. Louis, Mo.) per ml and incubated at 60°C for 1 h. After incubation, the sample was heated at 100°C for 5 min to denature the enzyme and centrifuged at 4,500 \times g for 15 min to collect the cell debris. The resultant pellet was resuspended in PBS to an A_{550} of 0.9 prior to use in the lectin typing assay.

Lectin typing. The freeze-dried native lectins from the following organisms were purchased from Sigma Chemical Co.: *Erythrina cristigali* (lectin ECA), *Lotus tetragonolobus* (lectin LotusA), *Triticum vulgare* (lectin WGA), and *Ulex europaeus* (lectin UEA-I). The nominal sugar specificities of these lectins are as follows: ECA, galactose; LotusA, fucose; WGA, *N*-acetylglucosamine; and UEA-I, fucose. The native lectins were each dissolved in PBS containing 0.02%

CaCl₂ and 0.02% MgCl₂ at a concentration of 0.5 mg/ml (10). Bacterial samples (40 μ l) were mixed with 10 μ l of lectin solution (0.5 mg/ml) in U-shaped microtiter wells (Bibby Sterilin Ltd., Stone, United Kingdom) for 5 s or, alternatively, were mixed with 10 μ l of PBS as a negative control and allowed to settle overnight, undisturbed, at 20°C. Results were read by visual inspection, whereby the formation of a carpet of aggregated cells on the bottoms of the wells indicated a positive reaction, whereas the gathering of cellular material into a dot on the bottoms of the wells indicated a negative result. Negative results were confirmed by tilting the wells at an angle greater than 45° and observing the movement of cellular material along the bottom of the well. As a positive control for lectin binding, individual lectins were shown to agglutinate a 0.75% (vol/vol) suspension of human type O erythrocytes after incubation at 20°C for 2 h in U-shaped microtiter wells.

SDS-PAGE analysis of LPS. Selected samples of bacterial biomass prepared for lectin typing were analyzed by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) with the discontinuous buffer system described previously (15). Electrophoresis was conducted with a loading gel of 5% acrylamide, a running gel of 15% acrylamide containing 3.2 M urea, and a constant current of 35 mA. Following electrophoresis, the gels were fixed and LPS was detected by silver staining (30).

Statistical analysis. The two-tailed Fisher exact test was used to compare the distributions of the lectin reaction patterns between characteristics. A *P* value of <0.05 was considered significant.

RESULTS

Reproducibility. All strains were lectin typed at the Department of Microbiology, National University of Ireland, Galway, which was the centralized facility. In addition, two independent subsets of these strains, 116 (38%) and 19 (6%) strains, respectively, were retested at two other test centers in Sweden (Department of Medical Microbiology, Dermatology and Infection, Lund University) and Estonia (Department of Medical Microbiology, University of Tartu) and yielded lectin reaction patterns identical to those observed at the centralized facility. Lectin typing of each extract was repeated in triplicate, and, in addition, strains that had been regrown from stocks were lectin typed again and gave identical results. The method proved to be robust and was readily applied to all the *H. pylori* isolates and in all the test laboratories. Furthermore, at the centralized facility, pellets of biomass that had been received and subcultures of frozen stocks gave identical results (data not shown).

Lectin reaction patterns. By using a panel of four lectins, a total of 16 distinct lectin reaction patterns are possible, and these were given arbitrary designations (Table 1) similar to those used in a previous study (10). Examples of all 16 possible reaction patterns were observed. The most common lectin reaction pattern detected was MH1, but other common patterns were MH5, MH10, MH13, and MH16 (Table 2). Isolates with pattern MH1 reacted with all the lectins in the panel, isolates with patterns MH5 and MH10 reacted with three of the four lectins, and isolates with pattern MH13 reacted with two of the four lectins, whereas isolates with pattern MH16 did not react with any of these lectins (Table 1). When excess biomass was available, representative strains with each of the lectin reaction patterns, including MH16, were analyzed by SDS-PAGE and were found to express high-molecular-weight LPS bearing an O side chain (data not shown).

Lectin reaction pattern and geographical variation. For meaningful analysis, strains were grouped into regions on the basis of their country of isolation. Five regions were delineated: Northern Europe comprised Estonia, Finland, and Sweden; Central Europe comprised Belgium, France, Germany, and Ireland; Eastern Europe comprised Bulgaria, the Czech

TABLE 1. Typing scheme^a for *H. pylori* based on reactions with four lectins

Lectin reaction pattern	Result of test with:			
	ECA	LotusA	UEA-I	WGA
MH1	+	+	+	+
MH2	+	+	+	-
MH3	+	+	-	-
MH4	+	-	-	-
MH5	-	+	+	+
MH6	-	-	+	+
MH7	-	-	-	+
MH8	+	-	-	+
MH9	+	-	+	+
MH10	+	+	-	+
MH11	-	+	-	-
MH12	-	-	+	-
MH13	-	+	-	+
MH14	+	-	+	-
MH15	-	+	+	-
MH16	-	-	-	-

^a Modified scheme from Hynes et al. (10).

Republic, Hungary, and Poland; Southern Europe comprised Greece, Italy, Portugal, and Spain; and the Pacific region (Asia and Oceania) comprised China, Japan, and the Polynesian population of New Zealand.

The occurrence of the 16 possible lectin reaction patterns in the different regions is shown in Table 2. Isolates from the Pacific region and Eastern Europe had a notably smaller variety of reaction patterns than those from Central, Southern, or Northern Europe. In particular, pattern MH1 was the most common pattern among Pacific region isolates, but it was also the most common among isolates from Central, Southern, and Eastern Europe. In contrast to the other regions, pattern MH10 occurred as frequently as pattern MH1 among Northern European isolates. Pattern MH16, isolates of which are nonreactive with any lectins in the panel, also was common

TABLE 2. Lectin reaction patterns of 309 *H. pylori* isolates from Pacific and European populations

Lectin reaction pattern	No. (%) of isolates					Total
	Pacific region	Europe				
		Northern	Central	Southern	Eastern	
MH1	22	24	16	36	20	118 (38.2)
MH2	1	1	2	1	0	5 (1.6)
MH3	0	1	1	1	0	3 (1.0)
MH4	0	2	2	1	0	5 (1.6)
MH5	5	13	7	5	3	33 (10.7)
MH6	0	0	1	3	0	4 (1.3)
MH7	3	4	2	1	0	10 (3.2)
MH8	2	6	1	2	2	13 (4.2)
MH9	0	0	0	0	3	3 (1.0)
MH10	5	24	1	7	8	45 (14.6)
MH11	0	2	3	1	0	6 (1.9)
MH12	0	1	0	1	0	2 (0.6)
MH13	2	12	3	4	2	23 (7.4)
MH14	0	1	1	0	0	2 (0.6)
MH15	0	2	0	6	1	9 (2.9)
MH16	2	20	1	5	0	28 (9.1)
Total	42	113	41	74	39	309 (100)

TABLE 3. Lectin reaction patterns of 90 *H. pylori* isolates from the European population and the host's associated pathology

Lectin reaction pattern	No. (%) of isolates from patients with the following:	
	Chronic gastritis	ACG ± IM ^a
MH1	28 (41.2)	9 (40.9)
MH2	1 (1.5)	
MH3	1 (1.5)	1 (4.5)
MH4		5 ^b (22.7)
MH5	11 (16.2)	1 (4.5)
MH6	2 (2.9)	
MH7	2 (2.9)	2 (9.1)
MH8	2 (2.9)	
MH9	1 (1.5)	
MH10	7 (10.3)	1 (4.5)
MH11	3 (4.4)	
MH12		
MH13	5 (7.4)	2 (9.1)
MH14	1 (1.5)	
MH15	2 (2.9)	
MH16	2 (2.9)	1 (4.5)
Total	68 (100)	22 (100)

^a Both antral and fundic atrophic gastritis.

^b The result is considered statistically significant ($P < 0.05$) by Fisher's exact test.

among Northern European isolates, but none of the Eastern European isolates had pattern MH16. Pattern MH9 was specific for the Eastern European isolates and occurred only in isolates from Poland and Bulgaria. Examination of the distribution of the lectin reaction patterns among the individual countries of the Northern European region demonstrated that whereas pattern MH10 was the most common among Swedish isolates, pattern MH5 was the most common among Estonian isolates and pattern MH16 was present only in Swedish isolates. In contrast, the Finnish isolates examined had a distribution of reaction patterns similar to those of isolates from other European countries, with pattern MH1 being the most common pattern.

Lectin reaction pattern and associated pathology. The majority of the European isolates ($n = 267$) were from patients with either chronic gastritis or ACG ± IM alone. Quantitative histological data for isolates from 90 of these patients were available, and these were subjected to further analysis. Data were provided for both the antral and the fundic regions of the patients' stomachs and included parameters such as degree of inflammation, occurrence of ACG ± IM, surface epithelial damage, *H. pylori* colonization density, and production of lymphoid follicles. Further quantitative histological information was available for the duodenal region, including parameters similar to those described above: villous atrophy, chronic inflammation, acute inflammation, occurrence of antral IM, *H. pylori* colonization density, and lymphoid follicle production.

No lectin reaction pattern statistically correlated with any of the histological findings listed above for the antral or fundic regions or the duodenal regions of *H. pylori*-infected patients. Analysis of European strains was limited to comparisons between the distributions of the lectin reaction patterns of isolates from chronic gastritis patients and patients with ACG ± IM (Table 3). Although only five isolates were typed as MH4,

TABLE 4. Lectin reaction patterns of 76 *H. pylori* isolates from the Swedish population and the host's associated pathology

Lectin reaction pattern	No. (%) of isolates from patients with the following:		
	Nonulcer disease	Ulcer-associated disease	Gastric adenocarcinoma
MH1	4 (8.9)	8 ^a (29.6)	
MH2	1 (2.2)		
MH3	1 (2.2)		
MH4		2 (7.4)	
MH5		3 (11.1)	
MH6			
MH7	1 (2.2)	1 (3.7)	
MH8	2 (4.4)	2 (7.4)	
MH9			
MH10	4 (8.9)	11 ^a (40.7)	
MH11	3 (6.7)		
MH12			
MH13	12 ^a (26.7)		
MH14			
MH15	1 (2.2)		
MH16	16 ^a (35.6)		4
Total	45 (100)	27 (100)	

^a The result is considered significant ($P < 0.05$) by Fisher's exact test.

a relationship was observed between isolates with lectin reaction pattern MH4 and ACG \pm IM, in which isolates with lectin reaction pattern MH4 were isolated from patients with ACG \pm IM, whereas isolates with this pattern were not isolated from patients with chronic gastritis ($P = 0.0006$) (Table 3). In addition, a relatively large proportion (92%) of isolates with pattern MH5 were associated with chronic gastritis, but this was not statistically significant ($P = 0.28$) compared with the other lectin reaction patterns detected in association with chronic gastritis and ACG \pm IM.

The Swedish isolates examined in the present study were isolated from patients with dyspeptic symptoms presenting for upper gastrointestinal endoscopy at an open-access endoscopy unit. As shown in Table 4, strains were selected on the basis of their availability from patients with gastritis (nonulcer disease), ulcerative disease, and gastric adenocarcinoma. Within this group of Swedish patients, isolates with lectin reaction pattern MH1 occurred significantly more often ($P = 0.047$) in those with ulcerative disease than in those with nonulcerative disease. Moreover, a stronger statistical relationship ($P = 0.002$) existed between pattern MH10 and ulcerative disease (Table 4). In contrast, lectin reaction patterns MH13 and MH16 occurred significantly more often ($P = 0.002$ and $P = 0.0002$, respectively) in isolates from Swedish patients with nonulcerative diseases than in isolates from Swedish patients with ulcerative diseases. Moreover, all four isolates from patients with adenocarcinoma, not defined as either an ulcer or a nonulcer disease, exhibited pattern MH16, isolates of which are unreactive with any of the lectins in the panel.

DISCUSSION

It has been suggested that the expression of carbohydrates in the O-polysaccharide side chain of *H. pylori* LPS, especially the mimicry of human Le antigens, plays a number of potential roles in the pathogenesis of the bacterium (22). They may act

in the evasion of host defenses, and subsequently, due to the chronic nature of the infection, their expression may play a role in autoimmune aspects of pathology (22). Also, it has been suggested that these antigens act in adhesion of the bacterium to the gastric mucosa (2, 7, 21). Furthermore, it has been reported that the expression of these fucosylated blood group antigens by *H. pylori* is related to the host Le phenotype (33), suggesting that carbohydrate expression by the bacteria adapts to the host environment. However, conflicting results have also been reported (8, 28).

The results of the present study indicate that considerable microdiversity exists in carbohydrate expression in the O side chain and LPS of *H. pylori*. In previous studies with antibodies, Mills et al. (19) developed a serotyping system using passive hemagglutination but could differentiate only six serotypes. Also, the extraction procedure used to produce the antigenic material and the technique used to titrate the antisera were time-consuming and labor-intensive, and the investigators concluded that they were unsuitable for use in the clinical laboratory. Nevertheless, in related immunoblotting studies, Mills et al. (19) demonstrated variations in the reactions of LPSs, particularly the O side chains, from isolates of various serotypes with their typing antisera. Furthermore, Simoons-Smit et al. (26) used four anti-Le^x and anti-Le^y monoclonal antibodies with different fine specificities in an enzyme-linked immunosorbent assay to differentiate strains into 12 different serotypes. However, a large proportion of isolates (15%) remained untypeable with anti-Le antibodies. In contrast to monoclonal antibodies that have been used to detect Le antigens (8, 11, 26, 32), lectins do not require complete epitopes for binding and may therefore detect more subtle differences in carbohydrate structure (5, 34). Previously, we have suggested that the *Anguilla anguilla* lectin, which has a specificity for fucose monosaccharide, may be used to subtype certain lectin reaction patterns (10). However, in the present study the ubiquitous reactivity of this lectin with all isolates examined (data not shown) meant that it was of little use for differentiation of strains. By the use of four lectins, all 16 possible lectin reaction patterns were observed in the course of the study, illustrating the presence of greater variation in carbohydrate expression by strains than that detectable with anti-Le monoclonal antibodies (8, 11, 26, 32).

The most common lectin reaction patterns observed were MH1, MH5, MH10, and MH13. All of these patterns exhibit reaction with at least LotusA and WGA, which have specificities for fucose and *N*-acetylglucosamine, respectively. However, the ability to differentiate between these isolates by the use of other lectins confirms the high degree of microdiversity in carbohydrate expression occurring in the LPS of *H. pylori*.

Also, the lectin reaction patterns of *H. pylori* isolates were analyzed in the context of their geographical region of isolation, and strain diversity was reflected in a number of region-specific variations in *H. pylori* lectin reaction patterns. No two regions or countries had the same frequency of distribution of patterns. Nevertheless, a region-specific reaction pattern was observed for isolates from Eastern Europe; pattern MH9 was observed only in isolates from Bulgaria and Poland. When data for the patients infected with these three isolates were more closely scrutinized, it was apparent that apart from their geographical origin, all three patients presented with common

pathologies and blood groups (acute gastritis, blood group A, and Rh factor positive). Although the number of isolates involved was small, it indicates the potential for use of lectin typing for the identification of geography-specific reaction patterns and may be indicative of the adaptation of strains to the host (4, 33). Greater variations in reaction patterns were associated with Northern, Central, and Southern European isolates. These greater variations in phenotypes may reflect the greater heterogeneity of these host populations. Conversely, isolates from populations with similar ethnic backgrounds, such as populations from China and Japan, had similar, but not identical, frequencies of lectin reaction patterns. Campbell et al. (4) showed that *H. pylori* strains isolated from the Polynesian and European descendant populations of New Zealand were genetically distinct. The MH1 pattern was the most common reaction pattern among isolates from the Pacific region, including Polynesian isolates, examined in the present study, as was the case for isolates from certain European populations; in contrast, the isolates from the Pacific region had relatively few lectin reaction patterns. These phenotypic variations of *H. pylori* which coincide with differences in human populations would argue for the presence of host-adapted strains (6). The presence of regional differences between *H. pylori* isolates observed in the present study is in agreement with the findings of a number of studies that have demonstrated genotypic differences between Asian and European isolates of *H. pylori* (12, 18, 31, 36). Moreover, Monteiro et al. (20) have shown in structural studies of LPS that although Le antigens are commonly expressed by strains from both the European and the Asian populations, there is a preponderance of Le^a and Le^b over Le^x and Le^y isoforms among Asian isolates. This agrees with data from the present study, which show a predominance of lectin reaction pattern MH1 among isolates from the Pacific region (Asia and Oceania), indicating the common occurrence of Le-related carbohydrate structures in this geographical region. Furthermore, similar to LPS structural studies, serological analyses with anti-Le antibodies have shown geographical differences in the expression of Le antigens by *H. pylori* strains, in particular between Asian and European isolates in the expression of Le^a and Le^b antigens (1, 8, 17, 26, 35, 36), although some isolates remain untypeable with anti-Le antibodies (8, 11, 26, 32). Importantly, the present lectin typing system has detected greater diversity among isolates than has been detectable with anti-Le antibodies (26) and has highlighted geographical differences in carbohydrate expression by *H. pylori* isolates due to its capacity for broader recognition of variations in carbohydrate structures (10).

In the present study, detailed histopathological data were available only for isolates from European (Eurohepygast Study Group) patients suffering from chronic gastritis and ACG ± IM. For this population of isolates we demonstrated that lectin reaction pattern MH4 was associated with ACG ± IM, whereas it was not associated with chronic gastritis alone. Although this relationship was significant ($P = 0.0006$) and pattern MH4 occurred only in isolates associated with ACG ± IM, only five isolates were typed as MH4. Therefore, to fully establish the association, a larger number of strains would need to be examined. On the other hand, analysis of isolates from a Swedish population with a wide spectrum of associated diseases demonstrated an association of isolates with reaction

patterns MH1 and MH10 with ulcer disease ($P = 0.047$ and $P = 0.002$, respectively), whereas it demonstrated an association of isolates with patterns MH13 and MH16 with nonulcerogenic disease ($P = 0.002$ and $P = 0.0002$, respectively). Moreover, isolates from patients suffering from gastric adenocarcinoma were all typed as MH16, although, again, the number so typed was small (i.e., four isolates). Nevertheless, it should be considered that the occurrence of gastric adenocarcinoma in the Swedish population is low and the small number of isolates in the present study reflects the low rate of occurrence of this pathology in that population. Isolates with the latter lectin reaction pattern are unreactive with any of the lectins in the panel, and this result may be indicative of the presence of atypical LPS structures. Biomass was not available from these strains for SDS-PAGE analysis, and they are therefore the focus of further investigations.

Nonetheless, if it is accepted that the associations described above do not reflect sampling bias, the data suggest that carbohydrate expression may influence the clinical outcome of an infection with *H. pylori*. Furthermore, this result is in agreement with data from other investigations (8, 17, 32) which have all associated Le antigen expression with an aggressive pathology, including ulceration. In addition, a Japanese study has shown that the expression of LPS by *H. pylori* with a low level of antigenicity may be associated with the development of gastric cancer (35) and that strains unreactive with anti-Le antibodies exist in this population (1). Similarly, this low level of antigenicity may be manifest in a low level of reactivity with lectins, as evidenced by the association of isolates with pattern MH16 (which do not react with any lectins) with gastric adenocarcinoma and the association of isolates with pattern MH4 (which react only with ECA lectin) with ACG ± IM. This association may be indicative of a link between isolate lectin reaction pattern and disease. In addition, since conditions in the stomach change upon the development of ACG ± IM and other bacteria can proliferate in this environment, the association may instead indicate that *H. pylori* strains with pattern MH4 could be better adapted to these new conditions.

Collectively, our data show that lectin typing is an effective method for epidemiological studies. It was successfully applied to a large and diverse group of *H. pylori* isolates at multiple centers, and the study proved its ease of application. Moreover, the system as applied was capable of demonstrating phenotypic differences between populations of *H. pylori* strains.

APPENDIX

Members of the Eurohepygast Study Group include the following: F. Mégraud (Bordeaux, France), C. O'Morain (Dublin, Ireland), P. Sipponen (Espoo, Finland), A. Price (London, United Kingdom), P. Malfertheiner (Magdeburg, Germany), N. Munoz (Lyon, France), F. Dabis (Bordeaux, France), N. Figura (Siena, Italy), M. Plebani (Padua, Italy), C. Sakarovitch (Bordeaux, France), F. Nachit (Bordeaux, France), I. M. Arenas (San Sebastian, Spain), A. Blasi (Catania, Italy), J. F. Bretagne (Rennes, France), S. Chaussade (Paris, France), J. C. Delchier (Créteil, France), J. D. De Korwin (Nancy, France), D. Duda (Wroclaw, Poland), J. Dzieniszewski (Warsaw, Poland), B. Fixa (Hradec Kralové, Czech Republic), X. Gisbert (Madrid, Spain), A. Goldis (Timisoara, Romania), J. Guerre,

(Paris, France), L. Hryniewiecki (Poznan, Poland), G. Inserra (Catania, Italy), A. L. Karvonen (Espoo, Finland), Z. Knapik (Wroclaw, Poland), H. Kolke (Tartu, Estonia), H. Lamouliatte (Bordeaux, France), A. Le Coustumier (Nancy, France), K. Linke (Poznan, Poland), J. Lonovics (Szeged, Hungary), H. Maaros (Tartu, Estonia), S. Michopoulos (Athens, Greece), J. Monés (Barcelona, Spain), J. G. Pajarés, (Madrid, Spain), L. Paradowski (Wroclaw, Poland), U. Peitz (Magdeburg, Germany), S. Pinto (Lisbon, Portugal), M. Quina (Lisbon, Portugal), G. Trautmann (Nancy, France), and K. Tchernev (Sofia, Bulgaria).

ACKNOWLEDGMENTS

This study was supported by grants from the Irish Health Research Board, the Federation of European Microbiological Societies, and the European Molecular Biology Organization (to A.P.M. and S.O.H.); the Swedish Research Council (grant 16×04723, to T.W.); and the Estonian Ministry of Education (grant 0418, to M.M.).

REFERENCES

- Amano, K.-I., S. Hayashi, T. Kubota, N. Fujii, and S.-I. Yokota. 1997. Reactivities of Lewis antigen monoclonal antibodies with lipopolysaccharides of *Helicobacter pylori* strains isolated from patients with gastroduodenal diseases in Japan. *Clin. Diagn. Lab. Immunol.* **4**:540–544.
- Appelmek, B. J., M. A. Monteiro, S. L. Martin, A. P. Moran, and C. M. J. E. Vandembroucke-Grauls. 2000. Why *Helicobacter pylori* has Lewis antigens. *Trends Microbiol.* **8**:565–570.
- Atherton, J. C., R. M. Peek, Jr., K. T. Tham, T. L. Cover, and M. J. Blaser. 1997. Clinical and pathological importance of heterogeneity in *vacA*, the vacuolating cytotoxin gene of *Helicobacter pylori*. *Gastroenterology* **112**:92–99.
- Campbell, S., A. Fraser, B. Hollis, J. Schmid, and P. W. O'Toole. 1997. Evidence for ethnic tropism of *Helicobacter pylori*. *Infect. Immun.* **65**:3708–3712.
- Chatterjee, B. P., A. K. Guha, R. Pal, and M. Bhattacharyya. 1989. Lectin typing of *Pseudomonas aeruginosa* strains of different serogroups, Habs and Fisher types. *Zentbl. Bakteriell. Parasitenkd. Infektkrankh. Hyg. Abt. 1 Orig.* **271**:364–371.
- Covacci, A., J. L. Telford, G. Del Giudice, J. Parsonnet, and R. Rappuoli. 1999. *Helicobacter pylori* virulence and genetic geography. *Science* **284**:1328–1333.
- Edwards, N. J., M. A. Monteiro, G. Faller, E. J. Walsh, A. P. Moran, I. S. Roberts, and N. J. High. 2000. Lewis X structures in the O antigen side-chain promote adhesion of *Helicobacter pylori* to the gastric epithelium. *Mol. Microbiol.* **35**:1530–1539.
- Heneghan, M. A., C. F. McCarthy, and A. P. Moran. 2000. Relationship of blood group determinants on *Helicobacter pylori* lipopolysaccharide with host Lewis phenotype and inflammatory response. *Infect. Immun.* **68**:937–941.
- Hunt, R. H. 1996. The role of *Helicobacter pylori* in pathogenesis: the spectrum of clinical outcomes. *Scand. J. Gastroenterol.* **31**(Suppl. 220):3–9.
- Hynes, S. O., S. Hirmo, T. Wadström, and A. P. Moran. 1999. Differentiation of *Helicobacter pylori* isolates based on lectin binding of cell extracts in an agglutination assay. *J. Clin. Microbiol.* **37**:1994–1998.
- Hynes, S. O., and A. P. Moran. 2000. Comparison of three serological methods for detection of Lewis antigens on the surface of *Helicobacter pylori*. *FEMS Microbiol. Lett.* **190**:67–72.
- Ito, Y., T. Azuma, S. Ito, H. Miyaji, M. Hirai, Y. Yamazaki, F. Sato, T. Kato, Y. Kohli, and M. Kuriyama. 1997. Analysis and typing of the *vacA* gene from *cagA*-positive strains of *Helicobacter pylori* isolated in Japan. *J. Clin. Microbiol.* **35**:1710–1714.
- Kellens, J. T. C., J. A. Jacobs, W. J. Peumans, and E. E. Stobberingh. 1994. Agglutination of "Streptococcus milleri" by lectins. *J. Med. Microbiol.* **41**:14–19.
- Kidd, M., J. C. Atherton, A. J. Lastovica, and J. A. Louw. 1999. Clustering of South African *Helicobacter pylori* isolates from peptic ulcer disease patients is demonstrated by repetitive extragenic palindromic-PCR fingerprinting. *Gut* **45**:499–502.
- Laemmli, U. K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature (London)* **227**:680–685.
- Marshall, B., H. Royce, D. I. Annear, C. S. Goodwin, J. W. Pearman, J. R. Warren, and J. A. Armstrong. 1984. Original isolation of *Campylobacter pyloridis* from human gastric mucosa. *Microbios Lett.* **25**:83–88.
- Marshall, D. G., S. O. Hynes, D. C. Coleman, C. A. O'Morain, C. J. Smyth, and A. P. Moran. 1999. Lack of a relationship between Lewis antigen expression and *cagA*, *CagA*, *vacA* and *VacA* status of Irish *Helicobacter pylori* isolates. *FEMS Immunol. Med. Microbiol.* **24**:79–90.
- Miehke, S., K. Kibler, J. G. Kim, N. Figura, S. M. Small, D. Y. Graham, and M. F. Go. 1996. Allelic variation in the *cagA* gene of *Helicobacter pylori* obtained from Korea compared to the United States. *Am. J. Gastroenterol.* **91**:1322–1325.
- Mills, S. D., L. A. Kurjanczyk, and J. L. Penner. 1992. Antigenicity of *Helicobacter pylori* lipopolysaccharides. *J. Clin. Microbiol.* **30**:3175–3180.
- Monteiro, M., P. Zheng, B. Ho, S. Yokota, K. Amano, Z. Pan, D. E. Berg, K. H. Chan, L. L. MacLean, and M. B. Perry. 2000. Expression of histoblood group antigens by lipopolysaccharides of *Helicobacter pylori* strains from Asian hosts: the propensity to express type 1 blood-group antigens. *Glycobiology* **10**:701–713.
- Moran, A. P., E. Sturegård, H. Sjunnesson, T. Wadström, and S. O. Hynes. 2000. The relationship between O-chain expression and colonisation ability of *Helicobacter pylori* in a mouse model. *FEMS Immunol. Med. Microbiol.* **29**:263–270.
- Moran, A. P. 1999. *Helicobacter pylori* lipopolysaccharide-mediated gastric and extragastric pathology. *J. Physiol. Pharmacol.* **50**:787–805.
- O'Sullivan, N., J. Benjamin, and M. B. Skirrow. 1990. Lectin typing of *Campylobacter* isolates. *J. Clin. Pathol.* **43**:957–960.
- Rudi, J., C. Kolb, M. Maiwald, I. Zuna, A. von Herbay, P. R. Galle, and W. Stremmel. 1997. Serum antibodies against *Helicobacter pylori* proteins *VacA* and *CagA* are associated with increased risk for gastric adenocarcinoma. *Dig. Dis. Sci.* **42**:1652–1659.
- Sillakivi, T., H. Aro, M. Ustav, M. Peetsalu, A. Peetsalu, and M. Mikelsaar. 2001. Diversity of *Helicobacter pylori* genotypes among Estonian and Russian patients with perforated peptic ulcer, living in Southern Estonia. *FEMS Microbiol. Lett.* **195**:29–33.
- Simoons-Smit, I. M., B. J. Appelmek, T. Verboom, R. Negrini, J. L. Penner, G. O. Aspinall, A. P. Moran, S. Fei Fei, S. Bi-Shan, W. Rudnicka, A. Savio, and J. De Graaff. 1996. Typing of *Helicobacter pylori* with monoclonal antibodies against Lewis antigens in lipopolysaccharide. *J. Clin. Microbiol.* **34**:2196–2200.
- Sipponen, P., M. Kekki, and M. Siurala. 1991. The Sydney System: epidemiology and natural history of chronic gastritis. *J. Gastroenterol. Hepatol.* **6**:244–251.
- Taylor, D. E., D. A. Rasko, R. Sherburne, C. Ho, and L. D. Jewell. 1998. Lack of correlation between Lewis antigen expression by *Helicobacter pylori* and gastric epithelial cells in infected patients. *Gastroenterology* **115**:1113–1122.
- Tham, K. T., R. M. Peek, Jr., J. C. Atherton, T. L. Cover, G. I. Perez-Perez, Y. Shyr, and M. J. Blaser. 2001. *Helicobacter pylori* genotypes, host factors, and gastric mucosal histopathology in peptic ulcer disease. *Hum. Pathol.* **32**:264–273.
- Tsai, C.-M., and C. E. Frasch. 1982. A sensitive silver stain for detecting lipopolysaccharides in polyacrylamide gels. *Anal. Biochem.* **119**:115–119.
- Van der Ende, A., Z.-J. Pan, A. Bart, R. W. M. Van der Hulst, M. Feller, S.-D. Xiao, G. N. J. Tytgat, and J. Dankert. 1998. *cagA*-positive *Helicobacter pylori* populations in China and The Netherlands are distinct. *Infect. Immun.* **66**:1822–1826.
- Wirth, H. P., M. Q. Yang, M. Karita, and M. J. Blaser. 1996. Expression of the human cell-surface glycoconjugates Lewis-X and Lewis-Y by *Helicobacter pylori* isolates is related to *cagA* status. *Infect. Immun.* **64**:4598–4605.
- Wirth, H. P., M. Yang, R. J. Peek, Jr., K. T. Tham, and M. J. Blaser. 1997. *Helicobacter pylori* Lewis expression is related to the host Lewis phenotype. *Gastroenterology* **113**:1091–1098.
- Wong, K. H., S. K. Skelton, and J. C. Feeley. 1985. Interaction of *Campylobacter jejuni* and *Campylobacter coli* with lectins and blood group antibodies. *J. Clin. Microbiol.* **22**:134–135.
- Yokota, S.-I., K.-I. Amano, Y. Shibata, M. Nakajima, M. Suzuki, S. Hayashi, N. Fujii, and T. Yokochi. 2000. Two distinct antigenic types of the polysaccharide chains of *Helicobacter pylori* lipopolysaccharides characterized by reactivity with sera from humans with natural infection. *Infect. Immun.* **68**:151–159.
- Zheng, P. Y., J. Hua, K. G. Yeoh, and B. Ho. 2000. Association of peptic ulcer with increased expression of Lewis antigens but not *cagA*, *iceA*, and *vacA* in *Helicobacter pylori* isolates in an Asian population. *Gut* **47**:18–22.